

Package ‘footprint’

October 13, 2022

Title Calculate Air Travel Emissions

Version 0.1

Description A handy tool to calculate carbon footprints from air travel based on three-letter International Air Transport Association (IATA) airport codes or latitude and longitude. footprint first calculates the great-circle distance between departure and arrival destinations. It then uses the Department of Environment, Food & Rural Affairs (DEFRA) greenhouse gas conversion factors for business air travel to estimate the carbon footprint. These conversion factors consider trip length, flight class (e.g. economy, business), and emissions metric (e.g. carbon dioxide equivalent, methane).

License CC0

URL <https://github.com/acircleda/footprint>

BugReports <https://github.com/acircleda/footprint/issues>

Depends R (>= 3.5.0)

Imports airports, dplyr, magrittr, rlang

Suggests devtools, knitr, rmarkdown, testthat (>= 2.1.0), tibble, covr

VignetteBuilder knitr

Encoding UTF-8

LazyData true

RoxygenNote 7.1.1

NeedsCompilation no

Author Anthony Schmidt [aut, cre] (<<https://orcid.org/0000-0003-4478-0638>>),
Kasia Kulma [aut]

Maintainer Anthony Schmidt <aschmi11@utk.edu>

Repository CRAN

Date/Publication 2021-01-06 10:40:02 UTC

R topics documented:

airport_footprint	2
latlong_footprint	3

airport_footprint	<i>Calculate flight emissions based on airport code pairs</i>
-------------------	---

Description

A function that calculates emissions per flight based on pairs of three-letter airport codes, flight classes, and emissions metrics. Emissions are returned in kilograms of the chosen metric.

Usage

```
airport_footprint(departure, arrival, flightClass = "Unknown", output = "co2e")
```

Arguments

departure	a character vector naming one or more three-letter IATA (International Air Transport Association) airport codes for outbound destination
arrival	a character vector naming one or more three-letter IATA (International Air Transport Association) airport codes for inbound destination
flightClass	a character vector naming one or more flight class categories. Must be of the following "Unknown" "Economy", "Economy+", "Business" or "First". If no argument is included, "Unknown" is the default and represents the average passenger.
output	a single character argument naming the emissions metric of the output. For metrics that include radiative forcing, one of <ul style="list-style-type: none"> • "co2e" (carbon dioxide equivalent with radiative forcing) - default • "co2" (carbon dioxide with radiative forcing) • "ch4" (methane with radiative forcing) • "n2o" (nitrous oxide with radiative forcing) • Metrics without radiative forcing: "co2e_norf", "co2_norf", "ch4_norf", or "n2o_norf".

Details

Distances between airports are based on the Haversine great-circle distance formula, which assumes a spherical earth. They are calculated using the `airportR` package. The carbon footprint estimates are derived from the Department for Environment, Food & Rural Affairs (UK) 2019 Greenhouse Gas Conversion Factors for Business Travel (air): <https://www.gov.uk/government/publications/greenhouse-gas-reporting-conversion-factors-2019>

Value

a numeric value expressed in kilograms of chosen metric

Examples

```
# Calculations based on individual flights
airport_footprint("LAX", "LHR")
airport_footprint("LAX", "LHR", "First")
airport_footprint("LAX", "LHR", "First", "ch4")
airport_footprint("LAX", "LHR", output = "ch4")

# Calculations based on a data frame of flights
library(dplyr)
library(tibble)

travel_data <- tribble(~name, ~from, ~to, ~class,
                      "Mike", "LAX", "PUS", "Economy",
                      "Will", "LGA", "LHR", "Economy+",
                      "Elle", "TYS", "TPA", "Business")

travel_data %>%
  rowwise() %>%
  mutate(emissions = airport_footprint(from, to,
                                       flightClass = class,
                                       output="co2e"))
```

latlong_footprint	<i>Calculate flight emissions based on longitude and latitude pairs</i>
-------------------	---

Description

A function that calculates emissions per flight based on longitude and latitude, flight classes, and emissions metrics. Emissions are returned in kilograms of the chosen metric.

Usage

```
latlong_footprint(
  departure_lat,
  departure_long,
  arrival_lat,
  arrival_long,
  flightClass = "Unknown",
  output = "co2e"
)
```

Arguments

`departure_lat` a numeric vector of one or more latitudes for departure location
`departure_long` a numeric vector of one or more longitudes for outbound location
`arrival_lat` a numeric vector of one or more latitudes for arrival location

<code>arrival_long</code>	a numeric vector of one or more longitudes for arrival location
<code>flightClass</code>	a character vector naming one or more flight class categories. Must be of the following "Unknown" "Economy", "Economy+", "Business" or "First". If no argument is included, "Unknown" is the default and represents the average passenger.
<code>output</code>	character emissions metric of the output. For metrics that include radiative forcing, one of <ul style="list-style-type: none"> • "co2e" (carbon dioxide equivalent with radiative forcing) - default • "co2" (carbon dioxide with radiative forcing) • "ch4" (methane with radiative forcing) • "n2o" (nitrous oxide with radiative forcing) • Metrics without radiative forcing: "co2e_norf", "co2_norf", "ch4_norf", or "n2o_norf". #

Details

Distances between latitude and longitude pairs are based on the Haversine great-circle distance formula, which assumes a spherical earth. The carbon footprint estimates are derived from the Department for Environment, Food & Rural Affairs (UK) 2019 Greenhouse Gas Conversion Factors for Business Travel (air): <https://www.gov.uk/government/publications/greenhouse-gas-reporting-conversion-factors-2019>

Value

a numeric value expressed in kilograms of chosen metric

Examples

```
# Calculations based on individual flights
latlong_footprint(34.052235, -118.243683, 35.179554, 129.075638)
latlong_footprint(34.052235, -118.243683, 35.179554, 129.075638, "First")
latlong_footprint(34.052235, -118.243683, 35.179554, 129.075638, "First", "ch4")
latlong_footprint(34.052235, -118.243683, 35.179554, 129.075638, output = "ch4")

# Calculations based on a data frame of flight
library(dplyr)
library(tibble)

travel_data <- tribble(~name, ~departure_lat, ~departure_long, ~arrival_lat, ~arrival_long,
  # Los Angeles -> Busan
  "Mike", 34.052235, -118.243683, 35.179554, 129.075638,
  # New York -> London
  "Will", 40.712776, -74.005974, 51.52, -0.10)

travel_data %>%
  rowwise() %>%
  mutate(emissions = latlong_footprint(departure_lat,
                                       departure_long,
                                       arrival_lat,
```

```
arrival_long,  
output="co2e"))
```

Index

`airport_footprint`, [2](#)

`latlong_footprint`, [3](#)