# Spearfish Sample Database


## *ABSTRACT*

The *spearfish* sample database is being distributed in order to provide users with a solid database on which to work in learning the tools of GRASS. This document provides some general information about the database and the map layers available.


With the release of GRASS 4.0, the GRASS development staff is pleased to announce that the sample data set *spearfish* is also being distributed. The *spearfish* data set covers two topographic 1:24,000 quads in western South Dakota. The names of the quads are Spearfish and Deadwood North, SD. The area covered by the data set is in the vicinity of Spearfish, SD and includes a majority of the Black Hills National Forest (i.e., Mount Rushmore). It is anticipated that enough data layers will be provided to allow users to use nearly all of the GRASS tools on the *spearfish* data set.


A majority of this *spearfish* database was initially provided to USA-CERL by the EROS Data Center (EDC) in Sioux Falls, SD. The GRASS Development staff expresses acknowledgement and thanks to the U.S. Geological Survey (USGS) and EROS Data Center for allowing us to distribute this data with our release of GRASS software. Thanks also to SPOT Image Corporation for providing multispectral and panchromatic satellite imagery for a portion of the *spearfish* data set and for allowing us to distribute this imagery with GRASS software. In addition to the data provided by the EDC and SPOT, researchers at USA-CERL have developed several new layers, thus enhancing the *spearfish* data set.


In order to use the *spearfish* data, when entering GRASS, enter *spearfish* as your choice for the current location.

```
- - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - - -

LOCATION: . . . . . .      spearfish_____________________

MAPSET: . . . . . .        your login-name_____________

_____________________________________________________________________
```

Available map layers for the *spearfish* database are detailed on the following page.

Raster files:*

| CELL MAP LAYERS | | | |
|---|---|---|---|
| FILENAME | TITLE | categories | bytes |
| aspect | Aspect | 25<br>15 degrees per category | 294978 |
| bugsites | Mountain Pine Beetle Damage | 1 | 26600 |
| density | Forest Density | 1-4 | 26600 |
| elevation.dem | Digital Elevation Model (7.5 minute) | 1066-1840 | 589956 |
| elevation.dted | Digital Terrain Elevation Data (DTED-1) | 1-241,243,245-251,254,255 | 26600 |
| erode.index | Erosion Index (ratio of K factor/T factor) | 12 | 10389 |
| fields | SCS Farm Fields | 63 | 2849 |
| geology | Geology | 9 | 26600 |
| landuse | Land Use | 8 | 26600 |
| owner | Ownership | 1-2 | 26600 |
| quads | Quads | 2 | 26600 |
| railroads | Railroads | 2 | 1191 |
| roads | Roads | 5 | 46583 |
| rstrct.areas | Restricted Areas | 4 | 26600 |
| rush.bnoise | Noise contours | 53-80 | 8537 |
| rushmore | Camp Rushmore | 1 | 1529 |
| slope | Slope (degrees) | 0-48,50,53-54,82-85,87-89 | 295388 |
| slope.7 | Slope (reclassed to percent rise) | 7 | 16 |
| soils | Soils | 54 | 75517 |
| soils.Kfactor | Soil K factors (surface layer) | 7 | 16 |
| soils.Tfactor | T soil erosion factor | 5 | 16 |
| soils.br.depth | Depth to Bedrock(inches) | 5 | 16 |
| soils.ph | PH for Soils | 5 | 16 |
| soils.range | Range Type | 12 | 16 |
| soils.texture | Soil Texture-USDA | 17 | 16 |
| spot.image | Spot multispectral band composite | | 1330000 |
| streams | Hydrography | 4 | 35200 |
| strm.dist | Proximity Analysis-Distance from Streams | 2 | 26600 |
| transport.misc | Miscellaneous Transportation Features | 2 | 1381 |
| trn.sites | Training Sites(Camp Rushmore) | 18 | 2701 |
| vegcover | Vegetation Cover | 1-6 | 26600 |

*In some map layers, there are gaps in the data and large areas of no data. Do not infer this to be a problem with the GRASS software. The source of the gaps and "no data" areas is within the map layer itself.

*"aspect" and "slope" were derived from "elevation.dem" using the program **slope.aspect**. They can be reclassed to suit the user. "slope.7" is a reclass of the "slope" layer containing seven categories of percent rise.

*"erode.index" was derived using r.mapcalc ((soils.Kfactor*100/soils.Tfactor)/100)

*Camp Rushmore is a **fictitious** installation created for demonstration purposes. "rushmore" is the boundary of this **fictitious** installation.

*"rush.bnoise" is example output from the program r.bnoise.

*In the file "slope", categories 83, 85, 88 and 89 actually represent areas in the NW and SE corners of the Spearfish database in which no elevation data exists. They appear as very steep slopes in the "slope" layer, because they are on the very edges of the data set; at the points where they occur, the data appears to drop off very sharply (to nothingness). In the file "slope.7" these categories were reclassed into the "no data" category.

*Mixed resolution within the data set reflects original data sources. The resolution can be changed, although higher resolutions will require more disk space.

DIG files:

| Layer name: | Layer description |
|---|---|
| fields | SCS farm fields |
| quads | USGS 1:24000 quads |
| railroads | Railroads |
| roads | Spearfish Roads |
| rstrct.areas | Restricted areas in Camp Rushmore |
| sections | Square mile sections |
| streams | Spearfish streams |
| t.9961.100 | Blockgroup 100 - Census tract 9961 |
| t.9961.100.all | All Tiger data - blockgroup 100 |
| t.9961.100.bks | All blocks - blockgroup 100 |
| t.9961.200 | Blockgroup 200 - Census tract 9961 |
| t.9961.200.all | All Tiger data - blockgroup 200 |
| t.9961.200.bks | All blocks - blockgroup 200 |
| t.9961.300 | Blockgroup 300 - Census tract 9961 |
| t.9961.300.all | All Tiger data - blockgroup 300 |
| t.9961.300.bks | All blocks - blockgroup 300 |
| t.9961.400 | Blockgroup 400 - Census tract 9961 |
| t.9961.400.all | All Tiger data - blockgroup 400 |
| t.9961.400.bks | All blocks - blockgroup 400 |
| t.9961.500 | Blockgroup 500 - Census tract 9961 |
| t.9961.500.all | All Tiger data - blockgroup 500 |
| t.9961.500.bks | All blocks - blockgroup 500 |
| t.county | Lawerance county boundary |
| t.hydro | Lawerance county hydrology |
| t.powerlines | Lawerance county powerlines |
| t.rails | Lawerance county railroads |
| t.roads | Lawerance county roads |
| t.roads.prime | Lawerance county primary highways |
| t.roads.second | Lawerance county secondary roads |
| t.tracts | Lawerance county census tracts |
| transport.misc | Miscellaneous transportation features |
| trn.sites | Training sites in Camp Rushmore |
| twp.range | Township and Range lines |

* rstrct.areas and trn.sites are **fictitious** DIG files for Camp Rushmore created for demonstration purposes.

*"streams" is a result of combining two seperate DLG files from Spearfish and Deadwood North quads, first by converting to GRASS vector files using **v.import**, and then patching the files together using **v.patch**.

*all vector files beginning with a 't.' prefix denote that they were derived from U.S. Bureau of the Census Tiger files. The TIGER file, developed by the Bureau of the Census, is a digital map of the entire United States and contains such features as roads, railroads, rivers, boundaries for census tracts and blocks, political areas such as cities and townships, feature names

and classification codes, alternate feature names, and within metropolitan areas, address ranges and ZIP Codes for streets.

The information contained within the TIGER file is unique within the United States and is broken down by state. Within each state, the TIGER information is further divided by county. For each county their are up twelve files or record types containing information relevant to that county.

Type 1 Lines with end nodes and basic attributes and geographic area codes
Type 2 Shape points between the end nodes
Type 3 Additional geographic area codes
Type 4 Additional feature identifier pointers
Type 5 Complete feature identifier list
Type 6 Additional address range and ZIP Code information
Type 7 Landmark features
Type 8 Area landmarks
Type I Area boundries
Type P Polygon locations
Type A Additional polygon geographic area codes
Type R Record number range

GRASS vector files derived from TIGER data were created using a database manager **RIM**, and the GRASS interface program, **v.db** (which alows interactive use of RIM in the GRASS environment). A RIM database was created for Lawerance County. Using **v.db**, different maplayers were then extracted out of the RIM database and converted into GRASS vector files.

The Spearfish data set is within Lawerance County, SD, and several maplayers for the county are included in the Spearfish mapset. All of the files listed below extend beyond the Spearfish DEFAULT region, and therefore, users should change their current geographic region to 'county' before viewing these files. 'county' is one of the GRASS geographic regions stored in the spearfish data set.

| | |
|---|---|
| t.county | Lawerance County boundary |
| t.hydro | All bodies of water within Lawerance County |
| t.powerlines | All power transmission lines within Lawerance County |
| t.rails | All railroad tracks within Lawerance County |
| t.roads | All roads contained within Lawerance County |
| t.roads.prime | All interstate roads within Lawerance County |
| t.roads.second | All U.S. Highways within Lawerance County |
| t.tracts | Census tract boundary outlines for tracts with the county |

In addition, several vector files are included for a sample census tract within Lawerance County. The tract, 9961, lies to the north and east of Spearfish and can be viewed after changing the current GRASS geographic region to '9961', which is also stored in the spearfish data set. The different '9961' vector files are break outs of the tract into three different formats. These files exist for census block-groups 100-500 within census tract 9961. The files and their naming schemes adhere to the following format.

| | |
|---|---|
| t.9961.100 | The outline of blockgroup 100 within tract 9961 |
| t.9961.100.all | Contains all of the TIGER data with blockgroup 100 |
| t.9961.100.bks | Outlines all of the blocks within the blockgroup 100 |

Imagery files:

| File name | File description |
|---|---|
| gs14.1 | Low Altitude Aerial Photograph |
| nhap.1 | NHAP blue raw band file |
| nhap.2 | NHAP green raw band file |
| nhap.3 | NHAP red raw band file |
| spot.comp | SPOT 3 band composite |
| spot.p | SPOT panchromatic band (10m) |
| spot.1.ms | Band 1- SPOT image (20m-green) |
| spot.2.ms | Band 2 - SPOT image (20m-red) |
| spot.3.ms | Band 3 - SPOT image (20m-near infrared) |

* Imagery files are found in LOCATION:imagery and MAPSET:spearfish, which is an x,y coordinate system.

* Imagery commands must be used to view these files. Please refer to the GRASS Imagery Tutorial.

*"gs14.1" is a low altitude aerial photograph which can be ortho-rectified using the the GRASS commands **i.mod.camera**, **i.build.blk**, and **i.rectify.blk**. Also included in this location is "gscam", a camera file containing information about the camera used to take the photograph. This file was created using **i.mod.camera**.

*NHAP and SPOT imagery are left as raw band files so that the user may georeference them to a desired coordinate system. The SPOT 3-band composite has been rectified to a UTM coordinate system and is located in LOCATION:spearfish and MAPSET:PERMANENT.

Site lists files:

| File name | File description | # of sites |
|---|---|---|
| archsites | Potential historic and archaeological sites | 25 |
| bugsites | Beetle Sites | 90 |

*"bugsites" has also been converted into a cell file. Analysis can only be done with cell files.

Three subregions for the data set are also provided. The name of the subregions are "county", "subregion.NW", and "9961". Users may change their current GRASS geographic region to one of these subregions within GRASS by using the GRASS command **g.region**.